



PERMIT TO ALLOW EMERGENCY USE OF AN AGVET CHEMICAL PRODUCT

PERMIT NUMBER -PER9375

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 29 SEPTEMBER 2006 31 DECEMBER 2006.

Permit Holder:

**DEPARTMENT OF AGRICULTURE AND FOOD WESTERN AUSTRALIA
3 Baron Hay Court
SOUTH PERTH WA 6151**

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Product to be used:

NUFARM FENITROTHION 1000 INSECTICIDE
DAVID GRAYS FENITROTHION 1000
RENTOKIL FENITROTHION 1000 INSECTICIDE
FARMOZ FENITROTHION 1000 INSECTICIDE
SUMITHION 1000EC INSECTICIDE
NEVWEB FENITROTHION 1000 GRAIN PROTECTANT
KENDON FENITROTHION 1000EC INSECTICIDE

Containing: 1000 g/L FENITROTHION as their only active constituent.

SUMITOMO SUMITHION ULV PREMIUM GRADE INSECTICIDE
Containing: 1.23 g/L FENITROTHION as the only active constituent.

Directions for Use:

| Crop | Pest | Rate |
|--|--------------------------|--|
| Failed crops of Canola, Cereals, Lupins, Faba beans, Peas, Chickpeas, Lentils and Vetch | Australian plague locust | EC products: 270 to 325mL per hectare ULV product: 200 to 400mL per hectare |

Critical Use Comments:

Apply in accordance with approved label for selected product.
Apply one application per failed crop only.
Use the higher rate through suitable boom spray in dense crops against adult locust.
DO NOT apply these products by aircraft at rates greater than 350mL per hectare.

Withholding Period:

Grazing: DO NOT graze for 7 days after application. Withhold livestock from slaughter for 14 days after application
Cutting for stock food: DO NOT cut for stock food for 14 days after application.

Where grazing stock have been oversprayed: DO NOT slaughter for 14 days after application.

EXPORT ANIMAL FEED INTERVALS:

WHERE STOCK ARE INTENDED FOR EXPORT IT SHOULD BE NOTED THAT **SAFEMEAT** HAVE RECOMMENDED A 14 DAY EXPORT ANIMAL FEED INTERVAL FOR THE USE OF FENITROTHION FOR PLAGUE LOCUST CONTROL. For further information contact Safemeat.

Jurisdiction:

WA only.

Additional Conditions:

THIS PERMIT provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

PERSONS who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit.

BUFFER ZONES:

Aerial application:

DO NOT apply at rates greater than 350mL/ha. DO NOT apply without a buffer zone AT LEAST 300m between the site of application and sensitive areas including (but not limited to) natural streams, rivers, wetlands, natural surface waters, waterways, human dwellings or neighbouring properties.

Ground application:

Apply fenitrothion by air-assisted, misting, electrostatic and boom sprayers.

Boom sprays:

DO NOT apply without a buffer zone of AT LEAST 50m where rates are 350mL/ha or below and AT LEAST 100m where rates are greater than 350mL/ha between the site of application and sensitive areas including (but not limited to) natural streams, rivers, wetlands, natural surface waters, waterways, human dwellings or neighbouring properties.

Misters:

DO NOT apply without a buffer zone of AT LEAST 100m at rates of 350mL/ha and below and AT LEAST 300m where rates are greater than 350mL/ha between the site of application and sensitive areas including (but not limited to) natural streams, rivers, wetlands, natural surface waters, waterways, human dwellings or neighbouring properties.

Issued by

Delegated Officer